arXiv:1502.00401v5 [math.AT] 15 Apr 2016

Topology of unitary groups and the prime orders of binomial coefficients

Haibao Duan*
Institute of Mathematics, Chinese Academy of Sciences,

Xianzu Lin†
College of Mathematics and Computer Science,
Fujian Normal University, Fuzhou, 350108, China;

April 18, 2016

Abstract

Let $c : SU(n) \to PSU(n) = SU(n)/\mathbb{Z}_n$ be the quotient map of the special unitary group $SU(n)$ by its center subgroup $\mathbb{Z}_n$. We determine the induced homomorphism $c^* : H^*(PSU(n)) \to H^*(SU(n))$ on cohomologies by computing with the prime orders of binomial coefficients.

2000 Mathematical Subject Classification: 55T10

Key words and phrases: Lie groups; cohomology, prime order of an integer

Email: dhb@math.ac.cn;linxianzu@126.com

1 Main result

The center subgroup of the special unitary group $SU(n)$ is the cyclic group $\mathbb{Z}_n$ of order n generated by the diagonal matrix $diag\{e^{\frac{2\pi}{n}i}, \cdots, e^{\frac{2\pi}{n}i}\} \in SU(n)$. Let $c : SU(n) \to PSU(n) := SU(n)/\mathbb{Z}_n$ be the quotient homomorphism. Consider the induced map on cohomologies

$$c^* : H^*(PSU(n); R) \to H^*(SU(n); R) \text{ with } R = \mathbb{Q} \text{ or } \mathbb{Z}.$$

It is well known that if $R = \mathbb{Q}$ then c^* is an isomorphism of algebras. It is also known that if $R = \mathbb{Z}$ there exist integral classes $\xi_{2r-1} \in H^{2r-1}(SU(n); \mathbb{Z})$, $2 \leq r \leq n$, so that the integral cohomology $H^*(SU(n); \mathbb{Z})$ is isomorphic to the exterior ring $\Lambda(\xi_3, \cdots, \xi_{2n-1})$. Conceivably, there exist integral cohomology classes $\zeta_{2r-1} \in H^{2r-1}(PSU(n); \mathbb{Z})$, as well as a sequence $(a_{n,2}, \cdots, a_{n,n})$ of integers, such that

---

[*] Supported by 973 Program 2011CB302400 and NSFC 11131008.

[†] Supported by National Natural Science Foundation for young (no.11401098).

(1.1) the set $\{1, \zeta_I = \underset{r\in I}{\Pi} \zeta_{2r-1} \mid I \subseteq \{2, \cdots, n\}\}$ of square free monomials is a basis of the free part of the ring $H^*(PSU(n); \mathbb{Z})$;

(1.2) the induced map of the covering c satisfies the relation

$c^*(\zeta_{2r-1}) = a_{n,r} \cdot \xi_{2r-1}$, $2 \leq r \leq n$.

See in the proof of Lemma 2.2 for the justification of the statements (1.1) and (1.2). In addition, since the mapping degree of c is the order of the center $\mathbb{Z}_n$, and since the product $\zeta_3 \cdots \zeta_{2n-1}$ is a generator of the top degree cohomology $H^m(PSU(n); \mathbb{Z}) = \mathbb{Z}$ ($m = \dim PSU(n)$) by (1.1), the calculation by (1.2)

$$c^*(\zeta_3 \cdots \zeta_{2n-1}) = (a_{n,2} \cdots a_{n,n}) \cdot \xi_3 \cdots \xi_{2n-1}$$

indicates that the sequence $(a_{n,2}, \cdots, a_{n,n})$ satisfies the constraint

(1.3) $n = a_{n,2} \cdot \cdots \cdot a_{n,n}$.

In view of (1.2) and (1.3) it is both of topological and arithmetic interests to study the problem of expressing the sequence $(a_{n,2}, \cdots, a_{n,n})$ by explicit function in n.

By *the prime factorization of an integer* $n \geq 2$ we mean the unique expression $n = p_1^{r_1} \cdots p_t^{r_t}$ with $1 < p_1 < \cdots < p_t$ the set of all prime factors of n. In term of this factorization one introduces the partition on the set $\{2, \cdots, n\}$ by

$$\{2, \cdots, n\} = Q_0(n) \underset{1\leq i\leq t}{\amalg} Q_{p_i}(n),$$

where $Q_{p_i}(n) = \{p_i^r \mid 1 \leq r \leq r_i\}$. The main result of this paper is

Theorem 1.1. *One has* $a_{n,k} = p_i$ *or* 1 *for* $k \in Q_{p_i}(n)$ *or* $k \in Q_0(n)$.

Theorem 1.1 implies that the decomposition (1.3) coincides with the prime factorization of the integer n. It plays also a role in computing the integral cohomology ring of the Lie group $PSU(n)$, compare Theorem 1.1 with [2, i), Lemma 4.6].

The remaining parts of the paper are so arranged. Based on a general approach to the cohomology theory of compact Lie groups developed in [2] an arithmetic characterization of the integers $a_{n,k}$ is obtained in Lemma 2.2. Section 3 constitutes the major contribution of the present work, where (granted with Lemma 2.2) Theorem 1.1 is established. In addition, in this paper the cohomologies are over the ring $\mathbb{Z}$ of integers, unless otherwise stated.

2 Preliminaries in topology

The center of the unitary group $U(n)$ of order n is the circle subgroup $S^1 = \{diag\{e^{i\theta}, \cdots, e^{i\theta}\} \mid \theta \in [0, 2\pi]\}$ that acts on $U(n)$ via right multiplication. The quotient group $PSU(n) := U(n)/S^1$ is the adjoint Lie group of the type A_{n-1}

[1], while the quotient map $C : U(n) \to PSU(n)$ is both a group homomorphism and an oriented circle bundle over $PSU(n)$. Let $\omega \in H^2(PSU(n))$ be the Euler class of the bundle C, and let $J_n(\omega) \subset H^*(PSU(n))$ be the subring generated by ω. For a pair (n, k) of integer with $1 \leq k \leq n$ we set

$$b_{n,k} = g.c.d.\{\tbinom{n}{1}, \cdots, \tbinom{n}{k}\},$$

where $\binom{n}{k} = \frac{n!}{k!(n-k)!}$ is the binomial coefficient.

Lemma 2.1. *The class ω generates the group $H^2(PSU(n)) = \mathbb{Z}_n$, and the subring $J_n(\omega)$ has the presentation*

$$(2.1)\quad J_n(\omega) = \mathbb{Z}[\omega] / \left\langle b_{n,k}\omega^k, 1 \leq k \leq n \right\rangle, \deg \omega = 2.$$

Proof. For a compact and 1–connected Lie group G whose center is the cyclic group $\mathbb{Z}_q$ of order q, consider the circle bundle over the quotient group $G/\mathbb{Z}_q$

$$C : \left[G \times S^1\right] / \mathbb{Z}_q \to G/\mathbb{Z}_q,$$

where $\mathbb{Z}_q$ acts on the circle group S^1 via the anti–clockwise rotation through angle $\frac{2\pi}{q}$. The first statement is verified by the facts that the Euler class of this bundle generates $H^2(G/\mathbb{Z}_q) = \mathbb{Z}_q$ (see [2, Theorem 1.3]), and that the unitary group $U(n)$ is isomorphic to $\left[SU(n) \times S^1\right] / \mathbb{Z}_n$.

Taking a maximal torus T on $PSU(n)$ the Borel transgression τ in the fibration $\pi : PSU(n) \to PSU(n)/T$ fits into the short exact sequence

$$(2.2)\quad 0 \to H^1(T) \xrightarrow{\tau} H^2(PSU(n)/T) \xrightarrow{\pi^*} H^2(PSU(n)) \to 0 \text{ (see [2, (2.5)]).}$$

Moreover, with respect to the canonical Schubert basis $\{\omega_1, \cdots, \omega_{n-1}\}$ on the cohomology $H^2(PSU(n)/T)$, the subgroup

$$\operatorname{Im} \tau = \ker \pi^* \subset H^2(PSU(n)/T)$$

is spanned by the subset (see [2, Theorem 2.4])

$$\{2\omega_1 - \omega_2, -\omega_{k-1} + 2\omega_k - \omega_{k+1}, -\omega_{n-2} + 2\omega_{n-1} \mid 2 \leq k \leq n-2\},$$

Consequently, the induced map π^* in (2.2) satisfies that

$$(2.3)\quad \pi^*(\omega_k) = k\omega,\ 1 \leq k \leq n-1.$$

On the other hand, by [2, i) of Theorem 5.1] the ring $H^*(PSU(n)/T)$ has the presentation

$$(2.4)\quad H^*(PSU(n)/T) = \mathbb{Z}[\omega_1, \cdots, \omega_{n-1}] / \left\langle c_r(\omega_1, \cdots, \omega_{n-1}), 2 \leq r \leq n \right\rangle$$

in which $c_r(\omega_1, \cdots, \omega_{n-1})$ is the r^{th} elementary symmetric polynomials in the set

$$\{\omega_1, \omega_k - \omega_{k-1}, -\omega_{n-1} \mid \ 2 \leq k \leq n-1\}$$

with cardinality n. By (2.3) and (2.4) the map π^* carries the ring $H^*(PSU(n)/T)$ surjectively onto the subring $J_n(\omega) \subset H^*(PSU(n))$, and induces the desired isomorphism

$$H^*(PSU(n)/T)/\left\langle \ker \pi^* \right\rangle = \mathbb{Z}[\omega_1]/\left\langle \tbinom{n}{r}\omega_1^r, 1 \leq r \leq n \right\rangle \cong J_n(\omega).\Box$$

Note that $b_{n,k} \mid b_{n,k-1}$ with $b_{n,1} = n$ and $b_{n,n} = 1$. The following result gives an arithmetic characterization for the integers $a_{n,k}$ in (1.2).

Lemma 2.2. $a_{n,k} = \frac{b_{n,k-1}}{b_{n,k}}$, $k \geq 2$.

Proof. The Gysin sequence of the circle bundle C provides us with an exact sequence relating the cohomologies $H^*(U(n))$ and $H^*(PSU(n))$

(2.5) $\cdots \to H^r(PSU(n)) \overset{C^*}{\to} H^r(U(n)) \overset{\theta}{\to} H^{r-1}(PSU(n)) \overset{\cup\omega}{\to} H^{r+1}(PSU(n)) \overset{C^*}{\to} \cdots,$

see [6, p.149] or [2, Theorem 1.3]. Let $\xi_{2r-1} \in H^{2r-1}(U(n))$ be the primary 1–form with characteristic polynomial $c_r(\omega_1, \cdots, \omega_{n-1})$ ([2, Definition 3.5]). Then

b) $\theta(\xi_{2k-1}) = \binom{n}{k}\omega^{k-1} \in J_n(\omega)$ (see [2, Formula (4.10)]).

In addition, in view of the canonical diffeomorphism $U(n) \cong S^1 \times SU(n)$ ([2, Theorem 1.2]) let $p : U(n) \to SU(n)$ be the projection onto the second factor. Then the induced map C^* factors through c^* in the fashion

c) $C^* = p^* \circ c^* : H^*(PSU(n)) \to H^*(SU(n)) \to H^*(U(n))$,

while the map p^* carries $H^*(SU(n))$ isomorphically onto the first summand of the decomposition by a)

d) $H^*(U(n)) = \Lambda(\xi_3, \cdots, \xi_{2n-1}) \oplus \xi_1 \cdot \Lambda(\xi_3, \cdots, \xi_{2n-1})$.

Comparing b) with the presentation (2.1) we find that the order of the class $\theta(\xi_{2k-1}) \in H^*(PSU(n))$ is precisely $\frac{b_{n,k-1}}{b_{n,k}}$. By the exactness of the sequence (2.5), and taking into account of the decomposition c), there exists a set

$$\Phi = \{\zeta_{2k-1} \in H^{2k-1}(PSU(n)) \mid 2 \leq k \leq n\}$$

of cohomology classes satisfying the relations

(2.6) $c^*(\zeta_{2k-1}) = \frac{b_{n,k-1}}{b_{n,k}} \cdot \xi_{2k-1}$, $2 \leq k \leq n$.

With (2.6) corresponding to (1.2) it suffices to show that the set $\{1, \zeta_I = \prod_{r \in I} \zeta_{2r-1} \mid I \subseteq \{2, \cdots, n\}\}$ of monomials satisfies the property (1.1).

Setting $m = \dim PSU(n)$ then the product $\xi_3 \cdots \xi_{2n-1}$ is a generator of the top degree cohomology group $H^m(SU(n)) = \mathbb{Z}$ by d). Since the mapping degree of c is n the calculation

$$c^*(\zeta_3 \cdots \zeta_{2n-1}) = b_{n,1} \cdot (\xi_3 \cdots \xi_{2n-1}) = n \cdot (\xi_3 \cdots \xi_{2n-1})$$

by (2.6) indicates that the monomial $\zeta_3 \cdots \zeta_{2n-1}$ is a generator of the top degree cohomology $H^m(PSU(n)) = \mathbb{Z}$. Consequently, by [3, Lemma 2.9] the set $\{1, \zeta_I = \underset{r\in I}{\Pi} \zeta_{2r-1} \mid I \subseteq \{2, \cdots, n\}\}$ of 2^{n-1} monomials is linearly independent, and spans a direct summand of the free part of the ring $H^*(PSU(n))$. The proof is now completed by the routine relation

$$\dim(H^*(PSU(n)) \otimes \mathbb{Q}) = 2^{n-1}\ .\square$$

Remark 2.3. In [1, §4] Baum and Browder obtained certain information on the Serre spectral sequence associated to the fiber sequence

$$U(n) \overset{C}{\to} PU(n) \overset{\chi}{\to} BS^1,$$

where χ is the classifying map of the Euler class $\omega \in H^2(PU(n))$. Precisely, write $H^*(U(n)) = \Lambda(z_i : i = 1, \cdots, n)$ and $H^*(BS^1) = \mathbb{Z}[\alpha]$, where $|z_i| = 2i - 1$ and $|\alpha| = 2$, Baum and Browder showed that the differentials d_r satisfy the relations

(2.7) $d_r(z_i) = 0$ for $r < 2i$; $d_{2i}(z_i) = \binom{n}{i}\alpha^i$.

From (2.7) one can obtain alternative proofs of Lemmas 2.1 and 2.2. For instance the subring $J_n(\omega)$ can be seen to be $\operatorname{Im}\chi = E_\infty^{*,0}(PU(n))$, see Ruiz [7].

In comparison our approach to Lemmas 2.2 is based on the exact sequence (2.5) which can be extended to compute the integral cohomologies of the adjoint Lie groups [2, Theorem 4.7, Theorem 4.12].$\square$

3 The prime orders of binomial coefficients

Let $\mathbb{R}^+$ and $\mathbb{N}$ be, respectively, the set of positive reals and the set of natural numbers. For a real $x \in \mathbb{R}$ let $[x]$ denote the unique integer satisfying $0 \leq x - [x] < 1$. It is straightforward to see that

Lemma 3.1. *For any* $x, y \in \mathbb{R}^+$, $m, n \in \mathbb{N}$, *with* $x + y = m + n$, *we have*

$$[x] + [y] \leq [m] + [n],$$

where the inequality holds if either $x \neq [x]$ *or* $y \neq [y]$.$\square$

Given a prime p and an $m \in \mathbb{N}$ *the order of* m *at* p, denoted by $ord_p(m)$, is the biggest integer a so that m is divisible by the power p^a. Clearly, if $n_1, \cdots, n_r$ is a sequence of positive integers then

(3.1) $ord_p(g.c.d.\{n_1, \cdots, n_r\}) = min\{ord_p(n_1), \cdots, ord_p(n_r)\}$.

The next formula is shown in [5, Theorem 416]

Lemma 3.2. $ord_p(n!) = \sum_{k=1}^{\infty}[\frac{n}{p^k}]$.

Applying Lemma 3.2 we show that

Lemma 3.3. *Assume that* $n = p^r n'$ *with* $(n\prime, p) = 1$*. Then* $k \in Q_p(n)$ *implies that the number* $a_{n,k}$ *is divisible by* p*.*

Proof. In view of the formula (3.1) and by Lemma 2.2, it suffices to show that

$$ord_p(\binom{n}{p^s}) < ord_p(\binom{n}{k})$$

for any $1 \leq k < p^s$ and $1 \leq s \leq r$. This is equivalent to

$$(3.2)\quad ord_p(p^s!(n-p^s)!) > ord_p(k!(n-k)!),\ 1 \leq k < p^s,\ 1 \leq s \leq r.$$

By Lemma 3.2 if $1 \leq k \leq p^s$ then

$$(3.3)\quad ord_p(k!(n-k)!) = \sum_{j=1}^{\infty}([\frac{n-k}{p^j}]+[\frac{k}{p^j}]) = \sum_{j=1}^{s}([\frac{n-k}{p^j}]+[\frac{k}{p^j}])+\sum_{j=s+1}^{\infty}[\frac{n-k}{p^j}],$$

where the second equality comes from the obvious relation

$$\sum_{j=s+1}^{\infty}[\frac{k}{p^j}] = 0.$$

Since the integer n is divisible by p^s the open interval $(\frac{n-p^s}{p^j}, \frac{n}{p^j})$ with $j > s$ contains no integer (for otherwise, there would be an integer $m \in (0, p^s)$ such that $n - m$ is divisible by the power p^j for some $j > s$, contradicting to that m is not divisible by p^s). It follows that

$$[\frac{n-p^s}{p^j}] = [\frac{n-k}{p^j}] \text{ for all } 1 \leq k < p^s,\ j > s.$$

Summing this equality over $j \geq s+1$ yields that

$$(3.4)\quad \sum_{j=s+1}^{\infty}[\frac{n-k}{p^j}] = \sum_{j=s+1}^{\infty}[\frac{n-p^s}{p^j}].$$

Finally, the relation (3.2) is shown by the calculation

$$\begin{aligned} ord_p(p^s!(n-p^s)!) &= \sum_{j=1}^{s}([\frac{n-p^s}{p^j}]+[\frac{p^s}{p^j}]) + \sum_{j=t+1}^{\infty}[\frac{n-p^s}{p^j}] \text{ (by (3.3))} \\ &> \sum_{j=1}^{s}([\frac{n-k}{p^j}]+[\frac{k}{p^j}]) + \sum_{j=s+1}^{\infty}[\frac{n-p^s}{p^j}] \text{ (by Lemma 3.1)} \\ &= \sum_{j=1}^{s}([\frac{n-k}{p^j}]+[\frac{k}{p^j}]) + \sum_{j=s+1}^{\infty}[\frac{n-k}{p^j}] \text{ (by (3.4))} \\ &= ord_p(k!(n-k)!) \text{ (by (3.3))}.\square \end{aligned}$$

Proof of Theorem 1.1. Assume that the prime factorization of the integer $n \geq 2$ is $n = p_1^{r_1} \cdots p_t^{r_t}$. For each $k \in Q_{p_i}(n)$ we can assume by Lemma 3.3 that

$$a_{n,k} = p_i \cdot \overline{a}_{n,k}$$

for some $\overline{a}_{n,k} \in \mathbb{N}$, and rephrase the decomposition (1.3) of the integer n as

$$\begin{aligned} n &= \prod_{k \in Q_{p_i}(n), 1 \leq i \leq t} a_{n,k} \prod_{k \in Q_0(n)} a_{n,k} \\ &= p_1^{r_1} \cdots p_t^{r_t} \prod_{k \in Q_{p_i}(n), 1 \leq i \leq t} \overline{a}_{n,k} \prod_{k \in Q_0(n)} a_{n,k}. \end{aligned}$$

Since the factor $p_1^{r_1} \cdots p_t^{r_t}$ on the right hand side coincides with the prime factorization of n, the equality above forces out the relations $\overline{a}_{n,k} = 1$ for $k \in Q_{p_i}(n)$, and $a_{n,k} = 1$ for $k \in Q_0(n)$. This completes the proof of Theorem 1.1.□

We conclude this paper with an application of Theorem 1.1. For a compact Lie group G with a maximal torus T let $\pi^* : H^*(G/T) \to H^*(G)$ be the induced map of the fibration $\pi : G \to G/T$. By Grothendieck [4] the subring $\operatorname{Im} \pi^* \subset H^*(G)$ is the *Chow ring* $A^*(G^c)$ of the reductive algebraic group G^c corresponding to G. From the proof of Lemma 2.1 we find that

(3.5) $A^*(PSU(n)^c) = J_n(\omega)$ (see (2.1)).

On the other hand, in view of the obvious decomposition $b_{n,k} = a_{n,k+1} \cdots a_{n,n}$ and by Theorem 1.1, the number $b_{n,k}$ admits the prime factorization

(3.6) $b_{n,k} = \prod_{1 \leq i \leq t} p_i^{r_i - l(i,k)}$,

where $l(i,k) = r_i$ if $k \geq p_i^{r_i}$, and satisfies the relation

$$p_i^{l(i,k)} \leq k < p_i^{l(i,k)+1}$$

if $k < p_i^{r_i}$. Combining (3.5) with (3.6) yields the following canonical decomposition of the ring $A^*(PSU(n)^c)$ into its primary torsion ideals

Corollary 3.4. *If $n > 2$ has the prime factorization $p_1^{r_1} \cdots p_t^{r_t}$, then*

$$A^*(PSU(n)^c) = \mathbb{Z} \underset{i \in \{1,\cdots,t\}}{\oplus} \mathbb{Z}[\omega]^+ / \left\langle p_i^{r_i}\omega, p_i^{r_i-1}\omega^{p_i}, p_i^{r_i-2}\omega^{p_i^2}, \cdots, \omega^{p_i^{r_i}} \right\rangle . \square$$

References

[1] P. F. Baum, W. Browder, The cohomology of quotients of classical groups, Topology 3 (1965), 305–336.

[2] H. Duan, Schubert calculus and cohomology of Lie groups. Part II. compact Lie groups, math.AT. arXiv:1502.00410.

[3] H. Duan, Xuezhi Zhao, Schubert calculus and cohomology of Lie groups. Part I. 1–connected Lie groups, math.AT (math.AG). arXiv:0711.2541.

[4] A. Grothendieck, Torsion homologique et sections rationnelles, Sem. C. Chevalley, ENS 1958, exposé 5, Secreatariat Math. IHP, Paris, 1958.

[5] G. Hardy and E. Wright, An introduction to the theory of numbers, Oxford Univ. Press, London, 1979.

[6] J. W. Milnor and J. D. Stasheff, Characteristic classes, Annals of Mathematics Studies, No. 76. Princeton University Press, 1974.

[7] C. A. Ruiz, The cohomology of the complex projective Stiefel manifold, Trans. Amer. Math. Soc. 146(1969), 541–547.